

Bacterial Assessment of the Cow's Milk from Different Areas in Baghdad City

Ali S. Hasan¹, Raad J. Hammadi², Wathiq Ali Hasson³ and Anas A. Humadi⁴

^{1,2} Department of Public Health, College of Veterinary Medicine, University of Dyala, Iraq.

³ Department of Public Health, College of Veterinary Medicine, University of Baghdad, Iraq.

⁴ Department of Pathology and poultry disease, College of Veterinary Medicine, University of Diyala, Iraq.

Corresponding author: Anas A. Humadi

Email : anashumady@yahoo.com

Abstract

The study was designed to investigate the raw cow's milk contamination with foodborne *Staphylococcus aureus* bacteria in different regions outside of Baghdad: (Mahmoudiyah, Al-Fadhilia, Abu-Ghraib, and Madinat Al-shuela). Milk samples (n=10) were randomly collected from healthy cows; media were prepared and cultured with diluted raw milk, the results of regions (Mahmoudiyah, Abu-Ghraib) showed that 1,0 out of 10 samples for both and 3, 2 of 10 samples were contaminated with *Staphylococcus aureus* bacteria of regions (Al-Fadhilia, Madinat Al-shuela). Therefore, in this study, we can estimate milk contamination with *Staphylococcus aureus* that may cause food poisoning if it is drunk by people. Therefore, we advise the manufacture of milk in high hygienic conditions and continuous pasteurization of milk, and treatment of the cases of subacute mastitis.

Keywords: Cow's milk, Staph. aureus, Baghdad city, Contamination.

INTRODUCTION

Food safety is a hot topic these days. Bacteria, fungi, viruses, and parasites are among the more than 200 diseases that food can spread. According to “public health and food safety” experts, foodborne germs cause millions of infections each year in the U.S.A. and across the globe. Yet, as stated by the “Centers for Disease Control and Prevention, the United States has one of the safest food sources in the world” (CDC, 2004).

The udder of the animal secretes milk as a sterile fluid. However, Contaminants can happen inside the udder and during the milking procedure, handling, and storage, in addition to the animals' diet, soil, and excrement being considered possible causes of contamination (Solomon *et al.*, 2013). Pathogenic bacteria should not be present in raw milk that is meant to be consumed by humans (Bertu *et al.*, 2010). Raw milk is thought to provide a perfect environment for aerobic bacteria, Lactococcus, Coliforms, Streptococcus, yeasts, and molds to develop and multiply (Solomon *et al.*, 2013). “Coliform bacteria” present in raw milk can be utilized as a sign of pathogenic bacteria contamination (Yuen *et al.*, 2012). Many initiatives are currently being made to use technologies to manufacture safe food supplies, such as low-additive dairy food in good quality "natural foods" (Leistner, 2002).

Raw milk is described as milk that was not been handled to change the quality or composition produced in compliance with recognized risky processing. "Milk products that have not been pasteurized or subjected to ultra-high temperature treatment" are defined as raw milk products.

Staphylococcus aureus contamination substantially impacts the safety and quality of dairy products. Because it produces staphylococcal enterotoxins, this bacterium is the most prevalent cause of food poisoning (SEs).

“*Staphylococcus aureus*” is one of the most prevalent bacteria that causes poisoning (Ioncarevic et al., 2005). It's a multipurpose human and animal pathogen that causes everything from minor skin infections to more serious ailments like pneumonia and septicemia (pelisser et al., 2009) (LOWY, 1998). Furthermore, human origins were found in 65 percent to 84 percent of all strains (Normanno *et al.*, 2007). Human skin, mucosa, nose, throat, hair, gastrointestinal, and urogenital tracts carry *S. aureus*. As a result, any product operated by hand is a staphylococcal potential source of infection. In addition, the risk of staphylococcal contamination of raw materials increases when suitable hygiene measures are not taken.

Staphylococcal food poisoning is a foodborne infection caused by staphylococcal enterotoxins generated in food by enterotoxigenic strains of *Staphylococcus aureus* (SFP). Milk is an excellent source of *S. aureus* development and enterotoxin production, and contaminating raw milk has been linked to several SFP outbreaks. (Asao *et al.*, 2003) Mastitis caused by *S. aureus* is a significant problem in the dairy industry, and diseased animals can affect bulk milk. Human workers, milking tools, the environment, or the udder and teat skin of dairy cows are all potential sources of bulk milk contamination; this is a problem in terms of food safety because enterotoxigenic *S. aureus* could cause SFP after consuming raw milk products (Jorgensen *et al.*, 2005).

S. aureus can manufacture a wide range of enterotoxins “(A, B, C, D, E, G, H, I, J, K, L, M, N, O, P, Q, R, and U),” It is usual for *S. aureus* to create these toxins at about the same time, despite the fact that classical enterotoxins are not produced by *S. aureus* “(A, B, C, D, E, G, H, I, J, K, L, M, N, O, P)” (Ietertre *et al.*, 2003).

“Food poisoning caused by *Staphylococcus aureus* (SFP)” is the third biggest foodborne illness globally even though it is a mild, self-limiting illness with a low fatality rate and is regarded as one of the most important economic illnesses (Cremonesi, 2005).

S. aureus bacteria can be killed by food processing without damaging SEs, which can withstand high temperatures and gastric proteases like pepsin (Cremonesi, 2007).

Due to contamination induced by inadequate hygienic standards or milk origin from mastitis cows, *S. aureus* is frequently discovered in raw milk and dairy products.

MATERIALS AND METHODS

Healthy calves were chosen at random to receive raw cow samples (n = 10) and taken to the laboratory in a refrigerator with an ice pack, in which they were treated one hour later.

Aseptic techniques were used to process the samples as soon as they arrived. “1mL of each milk sample was inoculated on Baird-Parker agar” to detect *S. aureus* (Difco, Detroit, Michigan, USA). Suspect colonies then “sub-cultured on blood agar plate (Difco, Detroit, Michigan, USA)” and incubated for twenty-four hours at 37 °C after 24 - 48 hours of incubation at 37 °C. (Jay, 2000) Gram stain, colony shape, catalase activity, “coagulase, and

Voges-Proskaver (VP0) tests were performed on suspicious colonies to identify *S. aureus* (Peles *et al.*, 2007).

RESULTS

S. aureus is typically found in raw “milk and dairy products” due to contamination caused by low sanitary standards or the origin of the milk, which can be from mastitic cows.

As noted in the below table, the percentage of infected raw milk samples of Mahmoudiyah was 10%, Al-Fadhilia 30%, Abu-Ghraib 0%, and Al-shuela 20%.

Table (1): The isolation percentages of raw milk samples contaminated by *S. aureus* that collected from districts of Baghdad province.

Region	Raw samples	Positive for <i>S. aureus</i>	Percentage
Mahmoudiyah	10	1	10%
Al-Fadhilia	=	3	30%
Abu-Ghraib	=	0	0%
Al-shuela	=	2	20%

DISCUSSION

REFERENCES:

1. Asao T., Kumeda Y., Kawai T., Shibata T., Oda H., Haruki K., Nakazawa H., Kozaki S.(2003). An extensive outbreak of staphylococcal food poisoning due to low-fat milk in Japan: estimation of enterotoxin A in the incriminated milk and powdered skim milk. *Epidemiol Infect.* 130: 33-40.
2. Asperger H. (1994). *Staphylococcus aureus*. In The significance of pathogenic microorganisms in raw milk. International Dairy Federation, Brussels. 24-42.
3. Bertu WJ, Dapar M, Gusi AM, Ngulukun SS, Leo S, Jwander LD. (2010). Prevalence of *brucella* antibodies in marketed milk in Jos and environs. *African Journal of Food Science.* 4(2):62-64.
4. Centers for Disease Control and Prevention. (CDC, 2004). Preliminary FoodNet data on the incidence of infection with pathogens transmitted commonly through food --- Selected Sites, United States, 2003 *MMWR.* 53:338.
5. Cremonesi P., Luzzana M., Brasca M., Morandi S., Lodi R., Vimercati C., Angellini D., Caramenti G., Moroni P., Castiglioni B. (2005). Development of a multiplex PCR assay for the identification of *Staphylococcus aureus* enterotoxigenic strains isolated from milk and dairy products. *Mol. Cell Probes* 19: 299-305.
6. Cremonesi P., Perez G., Pisoni G., Moroni P., Morandi S., Luzzana M., Brasca M., Castiglioni B.(2007). Detection of enterotoxigenic *Staphylococcus aureus* isolates in raw milk cheese. *Let. Appl. Microbiol.* 45: 586-91.
7. Desalegn A. (2014). Assessing the Microbiological quality of milk. *World Journal. Agron. Food Science. Technology.* 1(1): 1-10.
8. Jay JM. (2000). *Modern Food Microbiology* 6th ed. Aspen Publications Inc., Gaithersburg. Maryland, USA. 113-128.

9. Jorgensen H.J., Mork T., Caugant D.A., Kearns A., Rorvik L.M.(2005). Genetic variation among *Staphylococcus aureus* strains from Norwegian bulk milk. *Appl. Env. Microbiol.* 71: 8352-61.
10. Leistner L, Gould GW. *Hurdle technologies*(2002). Combination treatments for food stability, safety and quality. Kluwer/Plenum Publishers, New York, USA. 1-15.
11. Letertre, C., S. Perelle, F. dilasser, P. Fach (2003): Identifi cation of a new putative enterotoxin SEU encoded by the *egc* cluster of *Staphylococcus aureus*. *J. Appl. Microbiol.* 95: 38-43.
12. Loncarevic, S., H. J. jørgensen, A. Løvseth, T. Mathisen, L. M. Rørvik (2005). Diversity of *Staphylococcus aureus* enterotoxin types within single samples of raw milk and raw milk products. *J. Appl. Microbiol.* 98:344-50.
13. LOWY, F. D. (1998). *Staphylococcus aureus* infection. *N. Engl. J. Med.* 339: 520-32.
14. Normanno G., La Salandra G., Quaglia N.C., Corrente M., Parisi A., Santagada G., Firinu A., Crisetti E., Celano G.V. (2007): Occurrence, characterization and antimicrobial resistance of enterotoxigenic *Staphylococcus aureus* isolated from meat and dairy products. *International Journal of Microbiology.* 115: 290–96.
15. Peles, F., M. Wagner, L. Varga, I. Hein, P. Rieck, K. Gutser, P. Keresztúri, G.Kardos, I. Turcsányi, B. Béri, A. Szabó (2007). Characterization of *Staphylococcus aureus* strains isolated from bovine milk in Hungary. *Int. J. Food Microbiol.* 118: 186-93
16. Pelisser, M. R., C. S. Klein, K. R. Ascoli, T. R. Zotti, A. C. M. Arisil (2009). Occurrence of *Staphylococcus aureus* and multiplex PCR detection of classic enterotoxin genes in cheese and meat products. *Braz. J. Microbiol.* 40: 145-48.
17. Solomon M, Mulisa M, Yibeltal M, Desalegn G, Simenew K. (2013). Bacteriological quality of bovine raw milk at selected dairy farms in DebreZeit town, Ethiopia. *Comprehensive Journal of Food Science. Technol. Resarch.* 1(1):1-8.
18. Tortora GJ, Funke BR, Case CL. (2005). *Microbiological.* (8a edn), Porto Alegre, Artemed. 894.
19. Yuen SK, Yee CF, Yin FH. (2012). Microbiological quality and the impact of hygienic practices on the raw milk obtained, *international journal of Agri. and Food Science.* 2(2):55-59.