

“Mechanotransduction In Dental Implants”

Maaz Vohra¹, Nabeel Ahmed^{2*}, Khushali Shah³

¹Postgraduate Student, Department of Prosthodontics and Implantology Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai-600077, Tamil Nadu, India.
maazvohra390@gmail.com

^{2*}Associate Professor Department of Prosthodontics and Implantology Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai-600077, Tamil Nadu, India.
Email: nabeel.5610@gmail.com

³Assistant Professor Department of Prosthodontics and Implantology Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai-600077, Tamil Nadu, India.
Email: khushali.shah30@gmail.com

***Corresponding author:** Nabeel Ahmed

*Associate Professor Department of Prosthodontics and Implantology Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai-600077, Tamil Nadu, India.
Email: nabeel.5610@gmail.com

ABSTRACT

Dental implants made of titanium are placed directly into the bone tissue, which is a complex and dynamic tissue. This bone tissue not only helps maintain calcium homeostasis and serves as a hematopoietic organ, but it also acts as an immune regulator. The presence of a titanium implant during bone healing has recently been demonstrated to stimulate the immune system and cause type 2 inflammation, which appears to guide the connection between the host and the implant. This tends to show that osseointegration is a dynamic process involving a complicated series of reactions involving numerous host systems and pathways. Foreign Body Equilibrium (FBE) is a continuous equilibrium that occurs when osseointegration is not disrupted. To sustain the FBEs long-term integrity, the implant-host interaction must be improved further, notably when the mechanisms implicated in osseointegration breakdown start to function. The immune system may be stimulated as a result, disrupting the delicate balance between the osteoblast and the osteoclast, resulting in bone resorption. The immunomodulatory function of the Mesenchymal stem cell (MSC) has also been investigated, and in this vein, the notion of osseointegration immunomodulation through therapeutic mechanotransduction, particularly by extracorporeal shock waves therapy (ESWT), has recently been postulated. Mechanobiology has allowed us to investigate the impact of mechanical forces on cellular processes, revealing the intricate cellular control involved in mechanical signal transduction. Not only bone cells but also MSC, can be stimulated by mechanical stimulation. Mechanical stimulation can also alter the cellular shape of immune cells, affecting their phenotypic and function.

KEYWORDS: Mechanotransduction, Osteogenesis, BIC, Osteocytes, ESWT

INTRODUCTION

MECHANOTRANSDUCTION

We now know that osteogenesis is not solely dependent on the skeletal system's bone cells; rather, there is a multicellular collaboration. The complexity grows even more when we consider a fourth factor, mechanical stimuli, which can play a role in maintaining the material-host relationship's balance. Physical forces play a role in embryonic development, tissue homeostasis, and pathogenesis, among other things. The role of mechanical signals in controlling cellular activities, on the other hand, has just recently been acknowledged. As a result, novel study models to examine the influence of mechanical stresses on cells and tissues are being developed as interest in the topic of mechanobiology rises ⁹. Shear, tension, and compression forces all affect the cells. Tissues respond to mechanical signals in a variety of ways, including the creation of ECM components ¹⁰. Through the mechanotransduction signal, a mechanical change can influence gene expression and cellular activity ¹¹. The cytoskeleton filaments, such as actin and microtubules, would convey these mechanical impulses, which would then be converted into biochemical signals, with cell surface integrins crucial for mechanotransduction ¹². In fact, good clinical outcomes acquired by progressive loading protocol and instantaneous loading protocol ¹³ suggest that physiological mechanical stimuli can be advantageous to accompany the osseointegration of a dental implant and allow for successful osseointegration ¹⁴. In this regard, Duyck et al. demonstrated that mechanical stimulation can increase bone-implant contact (BIC) in an animal model using a bone chamber model ¹⁵. However, because mechanical stimuli are important for bone remodeling, further damage can result in implant failure ¹⁶. ESWT are commonly used in the area of therapeutic mechanotransduction at the moment. ESWT are supersonic waves that are created by a variety of technologies, including electrohydraulic, piezoelectric, electromechanical, and pneumatic devices, and cause transient pressure changes in the tissues they are applied to ¹⁷. The mechanical shocks created by ESWT are

thought to increase the permeability of the cell membrane, triggering the release of cytoplasmic ribonucleic acid (RNA) via an active process involving exosomes. This cellular event is the one that causes the effects seen in tissue repair to speed up. However, more research is needed to fully understand the underlying mechanism. The bone appears to be programmed to seal any portion that compromises its integrity, sealing and safeguarding the marrow content by restoring a cortical bone barrier. As a result, we might suppose that the "raw ingredients" (phosphate, calcium, and so on) are more readily available from a cortical bone source. In this regard, oral implants placed in low-density bone tissue (bone type IV) have been shown to have a greater failure rate¹⁸. On the other hand, there is an increase in peri-implant bone loss in patients with osteoporosis, despite the fact that there has been no difference observed yet in the survival rate of oral implants implanted in patients with and without osteoporosis¹⁹. In rats, Koolen et al. revealed at the histology level that after ESWT, *de novo* bone production occurs in bone defects restored with a titanium scaffold as a bone substitute. In a similar vein, Koolen et al. predicted that perioperative shock wave treatment could improve screw fixation and osseointegration of cortical and cancellous orthopedic screws, particularly in patients with osteoporosis. They were able to show that an ESWT performed immediately after the implantation of titanium screws (Ti6Al4V grade 5) increased screw attachment of the cortical screw, as measured by better mechanical strength and osseointegration, in a healthy rodent bone model. Shock waves have been suggested to cause the conversion of progenitor cells into osteogenic precursor cells, resulting in anabolic effects in the bone. Another theory is that ESWT causes osteocytic cell death via a process known as cavitation. This osteocyte death could stimulate local bone remodeling, prompting the osteoblast to make additional osteoid, and eventually leading to the formation of a neocortex²⁰. Osteocytes are mechanosensors and essential regulators of cellular homeostasis. The presence of HBMMSC and immune cells in the peri-implant milieu led us to believe that the ESWT may have an immunomodulatory effect favoring osseointegration. HBMMSCs can be identified not only in the peri-implant environment but also on the titanium surface. In monocytes, macrophages, and dendritic cells, mechanical signals play a significant role in the regulation of immunological and cellular processes²⁰. Immune cells that infiltrate the wound play a vital part in determining the wound's varying fate. The systemic depletion of macrophages, for example, has been shown to cause a persistent failure in the axolotl's regeneration potential, as well as significant fibrosis. However, after the macrophage population is restored, the axolotl's regeneration potential is restored²¹. While the use of immunomodulatory implants in and of themselves (clean implants) results in appropriate osseointegration, the FBE can be affected by clinical circumstances such as overload. ESWT is a promising therapeutic alternative to increase clinical success in oral implants and sustain FBE long-term due to the possibility of guiding the transition between the M1 inflammatory phase and the M2 anti-inflammatory phase by mechanotransduction. Because the capacity for tactile perception of osseointegrated implants, known as "osseoperception," grows over time²², this could potentially improve the feedback loop to the sensory brain. Furthermore, it has been suggested that adding nerve growth factor (NGF) to oral implants on a topical basis could aid to increase tactile sensitivity and reduce occlusal overload, and ESWT has been shown to be beneficial in enhancing NGF expression²³.

CONCLUSION

Mechanotransduction can help to strengthen the interaction between the implant and the host. However, research at the cellular and molecular level are required in order to determine the medical device as well as the most effective treatment range. All of this is being done in order to improve, maintain, and restore the balance of this trio of elements, namely bone cells, immune cells, and implants, which ultimately determines FBE's fate.

Conflict of Interest-None

REFERENCES

1. Trindade R, Albrektsson T, Wennerberg A. Current concepts for the biological basis of dental implants: foreign body equilibrium and osseointegration dynamics. *Oral Maxillofac Surg Clin North Am.* 2015 May;27(2):175–83.
2. Trindade R, Albrektsson T, Galli S, Prgomet Z, Tengvall P, Wennerberg A. Osseointegration and foreign body reaction: Titanium implants activate the immune system and suppress bone resorption during the first 4 weeks after implantation. *Clin Implant Dent Relat Res.* 2018 Feb;20(1):82–91.
3. Albrektsson T, Dahlin C, Jemt T, Sennerby L, Turri A, Wennerberg A. Is marginal bone loss around oral implants the result of a provoked foreign body reaction? *Clin Implant Dent Relat Res.* 2014 Apr;16(2):155–65.
4. Trindade R, Albrektsson T, Tengvall P, Wennerberg A. Foreign Body Reaction to Biomaterials: On Mechanisms for Buildup and Breakdown of Osseointegration. *Clin Implant Dent Relat Res.* 2016 Feb;18(1):192–203.
5. Albrektsson T, Canullo L, Cochran D, De Bruyn H. "Peri-Implantitis": A Complication of a Foreign Body or a Man-Made "Disease". Facts and Fiction. *Clin Implant Dent Relat Res.* 2016 Aug;18(4):840–9.
6. Nojehdehi S, Soudi S, Hesampour A, Rasouli S, Soleimani M, Hashemi SM. Immunomodulatory effects of mesenchymal stem cell-derived exosomes on experimental type-1 autoimmune diabetes. *J Cell Biochem.* 2018 Nov;119(11):9433–43.
7. Yusko EC, Asbury CL. Force is a signal that cells cannot ignore. *Mol Biol Cell [Internet].* 2017 Oct 13 [cited 2022 Jan 26]; Available from: <https://www.molbiolcell.org/doi/abs/10.1091/mbc.e13-12-0707>
8. Zhao L, Wu Z, Zhang Y. Low-magnitude mechanical vibration may be applied clinically to promote dental implant

- osseointegration. *Med Hypotheses*. 2009 Apr;72(4):451–2.
9. Guilak F, Cohen DM, Estes BT, Gimble JM, Liedtke W, Chen CS. Control of stem cell fate by physical interactions with the extracellular matrix. *Cell Stem Cell*. 2009 Jul 2;5(1):17–26.
 10. Leung DY, Glagov S, Mathews MB. Cyclic stretching stimulates synthesis of matrix components by arterial smooth muscle cells in vitro. *Science*. 1976 Feb 6;191(4226):475–7.
 11. Dunn SL, Olmedo ML. Mechanotransduction: Relevance to Physical Therapist Practice-Understanding Our Ability to Affect Genetic Expression Through Mechanical Forces. *Phys Ther*. 2016 May;96(5):712–21.
 12. MacKenna DA, Dolfi F, Vuori K, Ruoslahti E. Extracellular signal-regulated kinase and c-Jun NH2-terminal kinase activation by mechanical stretch is integrin-dependent and matrix-specific in rat cardiac fibroblasts. *J Clin Invest*. 1998 Jan 15;101(2):301–10.
 13. Piattelli A, Corigliano M, Scarano A, Costigliola G, Paolantonio M. Immediate loading of titanium plasma-sprayed implants: an histologic analysis in monkeys. *J Periodontol*. 1998 Mar;69(3):321–7.
 14. Sennerby L, Ericson LE, Thomsen P, Lekholm U, Astrand P. Structure of the bone-titanium interface in retrieved clinical oral implants. *Clin Oral Implants Res*. 1991 Jul;2(3):103–11.
 15. Badillo-Perona V, Cano-Sánchez J, Campo-Trapero J, Bascones-Martinez A. Peri-implant bone mechanobiology. Review of the literature. *Med Oral Patol Oral Cir Bucal*. 2011 Aug 1;16(5):e677–81.
 16. Albrektsson T, Chrcanovic B, Östman P-O, Sennerby L. Initial and long-term crestal bone responses to modern dental implants. *Periodontol 2000*. 2017 Feb;73(1):41–50.
 17. Ogden JA, Tóth-Kischkat A, Schultheiss R. Principles of shock wave therapy. *Clin Orthop Relat Res*. 2001 Jun;(387):8–17.
 18. Goiato MC, dos Santos DM, Santiago JF Jr, Moreno A, Pellizzer EP. Longevity of dental implants in type IV bone: a systematic review. *Int J Oral Maxillofac Surg*. 2014 Sep;43(9):1108–16.
 19. de Medeiros FCFL, Kudo GAH, Leme BG, Saraiva PP, Verri FR, Honório HM, et al. Dental implants in patients with osteoporosis: a systematic review with meta-analysis. *Int J Oral Maxillofac Surg*. 2018 Apr;47(4):480–91.
 20. Koolen MKE, Kruyt MC, Öner FC, Schaden W, Weinans H, van der Jagt OP. Effect of unfocused extracorporeal shockwave therapy on bone mineral content of twelve distal forearms of postmenopausal women: a clinical pilot study. *Arch Osteoporos* [Internet]. 2019 Nov 26 [cited 2022 Jan 26];14(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/31768768/>
 21. Godwin JW, Pinto AR, Rosenthal NA. Macrophages are required for adult salamander limb regeneration. *Proc Natl Acad Sci U S A*. 2013 Jun 4;110(23):9415–20.
 22. Mishra SK, Chowdhary R, Chrcanovic BR, Brånemark P-I. Osseoperception in Dental Implants: A Systematic Review. *J Prosthodont*. 2016 Apr;25(3):185–95.
 23. Lee J-H, Sung Y-B, Jang S-H. Nerve growth factor expression in stroke induced rats after shock wave. *J Phys Therapy Sci*. 2016 Dec;28(12):3451–3.